

# Implementation of Shi's non-degenerate Vuong test

June 8, 2025

## The original Vuong test

Vuong (1989) proposed a test for non-nested model. He considers two competing models,  $F_\beta = \{f(y|z; \beta); \beta \in \beta\}$  and  $G_\gamma = \{g(y|z; \gamma); \gamma \in \Gamma\}$ . Denoting  $h(y|z)$  the true conditional density. The distance of  $F_\beta$  from the true model is measured by the minimum KLIC:

$$D_f = E^0 [\ln h(y | z)] - E^0 [\ln f(y | z; \beta_*)]$$

where  $E^0$  is the expected value using the true joint distribution of  $(y, X)$  and  $\beta_*$  is the pseudo-true value of  $\beta$ <sup>1</sup>. As the true model is unobserved, denoting  $\theta^\top = (\beta^\top, \gamma^\top)$ , we consider the difference of the KLIC distance to the true model of model  $G_\gamma$  and model  $F_\beta$ :

$$\Lambda(\theta) = D_g - D_f = E^0 [\ln f(y | z; \beta_*)] - E^0 [\ln g(y | z; \gamma_*)] = E^0 \left[ \ln \frac{f(y | z; \beta_*)}{g(y | z; \gamma_*)} \right]$$

The null hypothesis is that the distance of the two models to the true models are equal or, equivalently, that:  $\Lambda = 0$ . The alternative hypothesis is either  $\Lambda > 0$ , which means that  $F_\beta$  is better than  $G_\gamma$  or  $\Lambda < 0$ , which means that  $G_\gamma$  is better than  $F_\beta$ . Denoting, for a given random sample of size  $N$ ,  $\hat{\beta}$  and  $\hat{\gamma}$  the maximum likelihood estimators of the two models and  $\ln L_f(\hat{\beta})$  and  $\ln L_g(\hat{\gamma})$  the maximum value of the log-likelihood functions of respectively  $F_\beta$  and  $G_\gamma$ ,  $\Lambda$  can be consistently estimated by:

$$\hat{\Lambda}_N = \frac{1}{N} \sum_{n=1}^N (\ln f(y_n | x_n, \hat{\beta}) - \ln g(y_n | x_n, \hat{\gamma})) = \frac{1}{N} (\ln L_f(\hat{\beta}) - \ln L_g(\hat{\gamma}))$$

---

<sup>1</sup> $\beta_*$  is called the pseudo-true value because  $f$  may be an uncorrect model.

which is the likelihood ratio divided by the sample size. Note that the statistic of the standard likelihood ratio test, suitable for nested models is  $2 \left( \ln L^f(\hat{\beta}) - \ln L^g(\hat{\gamma}) \right)$ , which is  $2N\hat{\Lambda}_N$ .

The variance of  $\Lambda$  is:

$$\omega_*^2 = V^o \left[ \ln \frac{f(y | x; \beta_*)}{g(y | x; \gamma_*)} \right]$$

which can be consistently estimated by:

$$\hat{\omega}_N^2 = \frac{1}{N} \sum_{n=1}^N \left( \ln f(y_n | x_n, \hat{\beta}) - \ln g(y_n | x_n, \hat{\gamma}) \right)^2 - \hat{\Lambda}_N^2$$

Three different cases should be considered:

- when the two models are nested,  $\omega_*^2$  is necessarily 0,
- when the two models are overlapping (which means than the models coincides for some values of the parameters),  $\omega_*^2$  *may be* equal to 0 or not,
- when the two models are stricly non-nested,  $\omega_*^2$  is necessarily strictly positive.

The distribution of the statistic depends on whether  $\omega_*^2$  is zero or positive. If  $\omega_*^2$  is positive, the statistic is  $\hat{T}_N = \sqrt{N} \frac{\hat{\Lambda}_N}{\hat{\omega}_N}$  and, under the null hypothesis that the two models are equivalent, follows a standard normal distribution. This is the case for two strictly non-nested models.

On the contrary, if  $\omega_*^2 = 0$ , the distribution is much more complicated. We need to define two matrices:  $A$  contains the expected values of the second derivates of  $\Lambda$ :

$$A(\theta_*) = E^0 \left[ \frac{\partial^2 \Lambda}{\partial \theta \partial \theta^\top} \right] = E^0 \left[ \begin{array}{cc} \frac{\partial^2 \ln f}{\partial \beta \partial \beta^\top} & 0 \\ 0 & -\frac{\partial^2 \ln g}{\partial \beta \partial \beta^\top} \end{array} \right] = \left[ \begin{array}{cc} A_f(\beta_*) & 0 \\ 0 & -A_g(\gamma_*) \end{array} \right]$$

and  $B$  the variance of its first derivatives:

$$B(\theta_*) = E^0 \left[ \frac{\partial \Lambda}{\partial \theta} \frac{\partial \Lambda}{\partial \theta^\top} \right] = E^0 \left[ \left( \frac{\partial \ln f}{\partial \beta}, -\frac{\partial \ln g}{\partial \gamma} \right) \left( \frac{\partial \ln f}{\partial \beta^\top}, -\frac{\partial \ln g}{\partial \gamma^\top} \right) \right] = E^0 \left[ \begin{array}{cc} \frac{\partial \ln f}{\partial \beta} \frac{\partial \ln f}{\partial \beta^\top} & -\frac{\partial \ln f}{\partial \beta} \frac{\partial \ln g}{\partial \gamma^\top} \\ -\frac{\partial \ln g}{\partial \gamma} \frac{\partial \ln f}{\partial \beta^\top} & \frac{\partial \ln g}{\partial \gamma} \frac{\partial \ln g}{\partial \gamma^\top} \end{array} \right]$$

or:

$$B(\theta_*) = \left[ \begin{array}{cc} B_f(\beta_*) & -B_{fg}(\beta_*, \gamma_*) \\ -B_{gf}(\beta_*, \gamma_*) & B_g(\gamma_*) \end{array} \right]$$

Then:

$$W(\theta_*) = B(\theta_*) [-A(\theta_*)]^{-1} = \begin{bmatrix} -B_f(\beta_*)A_f^{-1}(\beta_*) & -B_{fg}(\beta_*, \gamma_*)A_g^{-1}(\gamma_*) \\ B_{gf}(\gamma_*, \beta_*)A_f^{-1}(\beta_*) & B_g(\gamma_*)A_g^{-1}(\gamma_*) \end{bmatrix}$$

Denote  $\lambda_*$  the eigen values of  $W$ . When  $\omega_*^2 = 0$  (which is always the case for nested models), the statistic is the one used in the standard likelihood ratio test:  $2(\ln L_f - \ln L_g) = 2N\hat{\Lambda}_N$  which, under the null, follows a weighted  $\chi^2$  distribution with weights equal to  $\lambda_*$ . The Vuong test can be seen in this context as a more robust version of the standard likelihood ratio test, because it doesn't assume, under the null, that the larger model is correctly specified.

Note that, if the larger model is correctly specified, the information matrix equality implies that  $B_f(\theta_*) = A_f(\theta_*)$ . In this case, the two matrices on the diagonal of  $W$  reduce to  $-I_{K_f}$  and  $I_{K_g}$ , the trace of  $W$  to  $K_g - K_f$  and the distribution of the statistic under the null reduce to a  $\chi^2$  with  $K_g - K_f$  degrees of freedom.

The  $W$  matrix can be consistently estimate by computing the first and the second derivatives of the likelihood functions of the two models for  $\hat{\theta}$ . For example,

$$\hat{A}_f(\hat{\beta}) = \frac{1}{N} \sum_{n=1}^N \frac{\partial^2 \ln f}{\partial \beta \partial \beta^\top}(\hat{\beta}, x_n, y_n)$$

$$\hat{B}_{fg}(\hat{\theta}) = \frac{1}{N} \sum_{n=1}^N \frac{\partial \ln f}{\partial \beta}(\hat{\beta}, x_n, y_n) \frac{\partial \ln g}{\partial \gamma^\top}(\hat{\gamma}, x_n, y_n)$$

For the overlapping case, the test should be performed in two steps:

- the first step consists on testing whether  $\omega_*^*$  is 0 or not. This hypothesis is based on the statistic  $N\hat{\omega}^2$  which, under the null ( $\omega_*^2 = 0$ ) follows a weighted  $\chi^2$  distributions with weights equal to  $\lambda_*^2$ . If the null hypothesis is not rejected, the test stops at this step and the conclusion is that the two models are equivalent,
- if the null hypothesis is reject, the second step consists on applying the test for non-nested models previously described.

## The non-degenerate Vuong test

Shi (2015) proposed a non-degenerate version of the Vuong (1989) test. She shows that the Vuong test has size distortion, leading to subsequent overrejection. The cause of this problem is that the distribution of  $\hat{\Lambda}$  is discontinuous in the  $\omega^2$  parameter (namely a normal distribution if  $\omega^2 > 0$  and a distribution related to a weight  $\chi^2$  distribution if  $\omega^2 = 0$ ). Especially in small samples, it may be difficult to distinguish a positive versus a zero value of  $\omega^2$  because of sampling error. To solve this problem, using local asymptotic theory, Shi (2015) showed that, rewriting the Vuong statistic as:

$$\hat{T} = \frac{N\hat{\Lambda}_N}{\sqrt{N\hat{\omega}_N^2}}$$

the asymptotic distribution of the numerator and of the square of the denominator of the Vuong statistic is the same as:

$$\begin{pmatrix} N\hat{\Lambda}_N \\ N\hat{\omega}_N^2 \end{pmatrix} \rightarrow^d \begin{pmatrix} J_\Lambda \\ J_\omega \end{pmatrix} = \begin{pmatrix} \sigma z_\omega - z_\theta^\top V z_\theta / 2 \\ \sigma^2 - 2\sigma \rho_*^\top V z_\theta + z_\theta^\top V^2 z_\theta \end{pmatrix}$$

where:

$$\begin{pmatrix} z_\omega \\ z_\theta \end{pmatrix} \sim N \left( 0, \begin{pmatrix} 1 & \rho_*^\top \\ \rho_* & I \end{pmatrix} \right),$$

$\rho_*$  is a vector of length  $K_f + K_g$ ,  $\sigma$  a positive scalar and  $V$  is the diagonal matrix containing the eigen values of  $B^{\frac{1}{2}} A^{-1} B^{\frac{1}{2}}$ .

Based on this result, Shi (2015) showed:

- that the expected value of the numerator is  $-\text{trace}(V)/2$ , the classical Vuong statistic is therefore biased and this bias can be severe in small samples and when the degree of parametrization of the two models are very different<sup>2</sup>,
- that the denominator, being random, can take values close to zero with a significant probability, which can generate fat tails in the distribution of the statistic.

Shi (2015) therefore proposed to modify the numerator of the Vuong statistic:

$$\hat{\Lambda}_N^{\text{mod}} = \hat{\Lambda}_N + \frac{\text{tr}(V)}{2N}$$

and to add a constant to the denominator, so that:

$$\left( \hat{\omega}^{\text{mod}}(c) \right)^2 = \hat{\omega}^2 + c \text{tr}(V)^2 / N$$

The non-degenerate Vuong test is then:

$$T_N^{\text{mod}} = \frac{\hat{\Lambda}_N^{\text{mod}}}{\hat{\omega}^{\text{mod}}} = \sqrt{N} \frac{\hat{\Lambda}_N + \text{tr}(V)/2N}{\sqrt{\hat{\omega}^2 + c \text{tr}(V)^2 / N}}$$

---

<sup>2</sup>As the trace of  $V$  is the same as the trace of  $A^{-1}B$ , when the information matrix identity holds, it is equal to  $-K_f + K_g$ . The bias of the numerator is therefore caused by the difference in the degree of parametrization of the two models.

The distribution of the modified Vuong statistic can be estimated by simulations: drawing in the distribution of  $(z_\omega, z_\theta^\top)$ , we compute for every draw  $J_\Lambda$ ,  $J_\omega$  and  $J_\Lambda/\sqrt{J_\omega}$ . As  $\sigma$  and  $\rho_*$  can't be estimated consistently, the supremum over these parameters are taken, and Shi (2015) indicates that  $\rho_*$  should be in this case a vector where all the elements are zero except for the one that coincides with the highest absolute value of  $V$  which is set to 1.

The Shi test is then computed as follow:

1. start with a given size for the test, say  $\alpha = 0.05$ ,
2. for a given value of  $c$ , choose  $\sigma$  which maximize the simulated critical value for  $c$  and  $\alpha$ ,
3. adjust  $c$  so that this critical value equals the normal critical value, up to a small discrepancy (say 0.1); for example, if the size is 5%, the target is  $v_{1-\alpha/2} = 1.96 + 0.1 = 2.06$ ,
4. compute  $\hat{T}_N^{\text{mod}}$  for the given values of  $c$  and  $\sigma$  ; if  $\hat{T}_N^{\text{mod}} > v_{1-\alpha/2}$ , reject the null hypothesis at the  $\alpha$  level,
5. to get a p-value, if  $\hat{T}_N^{\text{mod}} > v_{1-\alpha/2}$  increase  $\alpha$  and repeat the previous steps until a new value of  $\alpha$  is obtained so that  $\hat{T}_N^{\text{mod}} = v_{1-\alpha^*/2}$ ,  $\alpha^*$  being the p-value of the test.

## Simulations

Shi (2015) provides an example of simulations of non-nested linear models that shows that the distribution of the Vuong statistic can be very different from a standard normal. The data generating process used for the simulations is:

$$y = 1 + \sum_{k=1}^{K_f} z_k^f + \sum_{k=1}^{K_g} z_k^g + \epsilon$$

where  $z^f$  is the set of  $K_f$  covariates that are used in the first model and  $z^g$  the set of  $K_g$  covariates used in the second model and  $\epsilon \sim N(0, 1 - a^2)$ .  $z_k^f \sim N(0, a/\sqrt{K_f})$  and  $z_k^g \sim N(0, a/\sqrt{K_g})$ , so that the explained variance explained by the two competing models is the same (equal to  $a^2$ ) and the null hypothesis of the Vuong test is true. The `vuong_sim` unables to simulate values of the Vuong test. As in Shi (2015), we use a very different degree of parametrization for the two models, with  $K_f = 15$  and  $K_G = 1$ .

```
library(miscr)
Vuong <- vuong_sim(N = 100, R = 1000, Kf = 15, Kg = 1, a = 0.5)
head(Vuong)
```

```
[1] 0.3544664 0.2017979 1.5595724 0.7240125 0.1898368 -0.2544294
```

```
mean(Vuong)
```

```
[1] 1.10883
```

```
mean(abs(Vuong) > 1.96)
```

```
[1] 0.214
```

We can see that the the mean of the statistic for the 1000 replications is far away from 0, which means that the numerator of the Vuong statistic is seriously biased. 21.4% of the values of the statistic are greater than the critical value so that the Vuong test will lead in such context a noticeable overrejection. The empirical pdf is shown in Figure 1, along with the normal pdf.




Figure 1: Empirical distribution of the Vuong statistic

## Implementation of the non-degenerate Vuong test

The `micsr` package provides a `ndvuong` function that implements the classical Vuong test. It has a `nest` argument (that is `FALSE` by default but can be set to `TRUE` to get the nested version

of the Vuong test). This package also provide a `llcont` generic which returns a vector of length  $N$  containing the contribution of every observation to the log-likelihood.

The `ndvuong` package provides the `ndvuong` function. As for the `vuongtest` function, the two main arguments are two fitted models (say `model1` and `model2`). The  $\hat{\Lambda}_n$  vector is obtained using `llcont(model1) - llcont(model2)`. The relevant matrices  $A_i$  and  $B_i$  are computed from the fitted models using the `estfun` and the `meat` functions from the sandwich package. More precisely,  $A^{-1}$  is `bdiag(-bread(model1), bread(model2))`<sup>3</sup> and  $B$  is `crossprod(estfun(model1), - estfun(model2)) / N`, where  $N$  is the sample size. Therefore, the `ndvuong` function can be used with any models for which a `llcont`, a `estfun` and a `bread` method is available.

## Applications

### Voter turnout

The first application is the example used in Shi (2015) and is used to compare our **R** program with Shi's **stata**'s program. Coate and Conlin (2004) used several models of electoral participation, using data concerning referenda about alcohol sales regulation in Texas. Three models are estimated: the preferred group-utilitarian model, a "simple, but plausible, alternative: the intensity model" and a reduced form model estimated by the seemingly unrelated residuals method. They are provided in the `ndvuong` package as `turnout`, a list of three fitted models<sup>4</sup>. The results of the Shi test are given below. We first compute the Shi statistic for an error level of 5%. We therefore set the `size` argument to 0.05 (this is actually the default value) and the `pval` argument to `FALSE`.

```
test <- ndvuong(turnout$group, turnout$intens, size = 0.05, pval = FALSE)
test
```

Non-degenerate Vuong test for non-nested models

```
data: turnout$group-turnout$intens
z = 1.7759, size = 0.050000, vuong_stat = 2.084528, constant =
0.381107, crit-value = 2.059963, sum e.v. = -10.997224, vuong_p.value =
0.018556
alternative hypothesis: different models
```

<sup>3</sup>`bdiag` if a function that construct a block-diagonal matrix from its arguments.

<sup>4</sup>The estimation is rather complicated because some linear constraints are used to compute the maximum likelihood estimator in Coate and Conlin (2004)'s **stata** script. This is the reason why we provide only the results of the estimations, performed using the `maxLik` package.

The Shi statistic is 1.776, which is smaller than the critical value 2.06. Therefore, based on the Shi test, we can't reject the hypothesis that the two competing models are equivalent at the 5% level. The value of the constant  $c$  is also reported, as is the sum of the eigen values of the  $V$  matrix (`sum e.v.`). The classical Vuong statistic is also reported (2.085) and is greater than the 5% normal critical value (the p-value is 0.019). Therefore, the classical Vuong test and the non-degenerate version lead to opposite conclusions at the 5% level.

To get only the classical Vuong test, the `nd` argument can be set to `FALSE`:

```
ndvuong(turnout$group, turnout$intens, nd = FALSE)
```

Vuong test for non-nested models

```
data: turnout$group-turnout$intens
z = 2.0845, p-value = 0.01856
alternative hypothesis: different models
```

To get the p-value of the non-degenerate Vuong test, the `pval` argument should be set to `TRUE`.

```
test <- ndvuong(turnout$group, turnout$intens, pval = TRUE)
test
```

Non-degenerate Vuong test for non-nested models

```
data: turnout$group-turnout$intens
z = 1.8125, vuong_stat = 2.084528, constant = 0.000000, sum e.v. =
-10.997224, vuong_p.value = 0.018556, p-value = 0.0864
alternative hypothesis: different models
```

The results indicate that the p-value is 0.086, which confirms that the Shi test concludes that the two model are equivalent at the 5% level.

## Transport mode choice (nested models)

The third example concerns transport mode choice in Canada. The dataset, provided by the `mlogit` package is called `ModeCanada` and has been used extensively in the transport demand literature (see in particular Bhat 1995; Koppelman and Wen 2000; and Wen and Koppelman

2001). The following example is from Croissant (2020). The raw data set is first transformed to make it suitable for the estimation of discrete choice models. The sample is restricted to the individuals for which 4 transport modes are available (bus, air, train and car).

```
if (requireNamespace("mlogit")){
  library(mlogit)
  data("ModeCanada", package = "mlogit")
  MC <- dfidx(ModeCanada, subset = noalt == 4)
}
```

We first estimate the simplest discrete choice model, which is the multinomial logit model. The bus share being negligible, the choice set is restricted to the three other modes and the reference mode is set to car.

```
if (requireNamespace("mlogit")){
  ml <- mlogit(choice ~ freq + cost + ivt + ovt | urban + income, MC,
    reflevel = 'car', alt.subset = c("car", "train", "air"))
}
```

This model relies on the hypothesis that the unobserved component of the utility functions for the different modes are independent and identical Gumbell variables. Bhat (1995) proposed the heteroscedastic logit for which the errors follow a general Gumbell distributions with a supplementary scale parameter to be estimated. As the overall scale of utility is not identified, the scale parameter of the reference alternative (car) is set to one.

```
if (requireNamespace("mlogit")){
  hl <- mlogit(choice ~ freq + cost + ivt + ovt | urban + income, MC,
    reflevel = 'car', alt.subset = c("car", "train", "air"),
    heterosc = TRUE)
  coef(summary(hl))
}
```

	Estimate	Std. Error	z-value	Pr(> z )
(Intercept):train	0.678393435	0.332762598	2.038671	4.148288e-02
(Intercept):air	0.656754399	0.468163091	1.402832	1.606668e-01
freq	0.063924677	0.004916769	13.001360	0.000000e+00
cost	-0.026961457	0.004283139	-6.294789	3.078178e-10
ivt	-0.009680773	0.001053874	-9.185892	0.000000e+00
ovt	-0.032165526	0.003593007	-8.952258	0.000000e+00
urban:train	0.797131578	0.120739176	6.602096	4.053868e-11
urban:air	0.445472634	0.082160945	5.421951	5.895197e-08
income:train	-0.012597857	0.003994180	-3.154053	1.610196e-03

```
income:air      0.018859983 0.003215926  5.864558 4.503294e-09
sp.train        1.237182865 0.110460959 11.200182 0.000000e+00
sp.air          0.540323852 0.111835294  4.831425 1.355592e-06
```

The two supplementary coefficients are `sp.train` and `sp.air`. The student statistics reported are irrelevant because they test the hypothesis that these parameters are 0, as the relevant hypothesis of homoscedasticity is that both of them equal one. The heteroscedastic logit being nested in the multinomial logit model, we can first use the three classical tests: the Wald test (based on the unconstrained model `hl`), the score test (based on the constrained model `ml`) and the likelihood ratio model (based on the comparison of both models).

To perform the Wald test, we use `lmtest::waldtest`, for which a special method is provided by the `mlogit` package. The arguments are the unconstrained model (`hl`) and the update that should be used in order to get the constrained model (`heterosc = FALSE`). To compute the scoretest, we use `mlogit::scoretest`, for which the arguments are the constrained model (`ml`) and the update that should be used in order to get the unconstrained model (`heterosc = TRUE`). Finally, the likelihood ratio test is performed using `lmtest::lrtest`.

```
if (requireNamespace("lmtest")){
  lmtest::waldtest(hl, heterosc = FALSE)
  scoretest(ml, heterosc = TRUE)
  lmtest::lrtest(hl, ml)
}
```

#### Likelihood ratio test

```
Model 1: choice ~ freq + cost + ivt + ovt | urban + income
Model 2: choice ~ freq + cost + ivt + ovt | urban + income
#Df  LogLik Df  Chisq Pr(>Chisq)
1  12 -1838.1
2  10 -1841.6 -2  6.8882    0.03193 *
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The three statistics are  $\chi^2$  with two degrees of freedom under the null hypothesis of homoscedasticity. The three tests reject the null hypothesis at the 5% level, and even at the 1% level for the Wald and the score test. These three tests relies on the hypothesis that, under the null, the constrained model is the true model. We can get rid of this hypothesis using a Vuong test. Note the use of the `nested` argument that is set to `TRUE`:

```
ndvuong(hl, ml, nested = TRUE)
```

### Non-degenerate Vuong test for nested models

```
data: hl-ml
z = 0.4554, vuong_stat = 6.888241, vuong_p.value = 0.047927, p-value =
0.211
alternative hypothesis: different models
```

The homoscedasticity hypothesis is still rejected at the 5% level for the classical Vuong test (the p-value is 0.048), but it is not using the non-degenerate Vuong test (p-value of 0.211).

### Transport mode choice (overlapping models)

We consider finally another dataset from **mlogit** called **RiskyTransport**, that has been used by León and Miguel (2017) and concerns the choice of one mode (among water-taxi, ferry, hovercraft and helicopter) for trips from Sierra Leone's international airport to downtown Freetown.

```
if (requireNamespace("mlogit")){
  library(mlogit)
  data("RiskyTransport", package = "mlogit")
  RT <- dfidx(RiskyTransport, idx = c(id = "chid", "mode"), choice = "choice")
}
```

We estimate models with only one covariate, the generalized cost of the mode. We estimate three models: the basic multinomial logit model, the heteroscedastic model and a nested model where alternatives are grouped in two nests according to the fact that they are fast or slow modes.

```
if (requireNamespace("mlogit")){
  ml <- mlogit(choice ~ cost, data = RT)
  hl <- mlogit(choice ~ cost, data = RT, heterosc = TRUE)
  nl <- mlogit(formula = choice ~ cost, data = RT,
              nests = list(fast = c("Helicopter", "Hovercraft"),
                           slow = c("WaterTaxi", "Ferry")),
              un.nest.el = TRUE)
}
```

Compared to the multinomial model, the heteroscedastic model has 3 supplementary coefficients (the scale parameters for 3 modes, the one for the reference mode being set to 1) and the nested logit model has one supplementary parameter which is the nest elasticity (iv in the table). Both models reduce to the multinomial logit model if:

- `sp.WaterTaxi = sp.Ferry = sp.Hovercraft = 1` for the heteroscedastic model,
- `iv = 1` for the nested logit model.

Therefore, the two models are over-lapping, as they reduce to the same model (the multinomial logit model) for some values of the parameters.

The first step of the test is the variance test. It can be performed using `ndvuong` by setting the argument `vartest` to `TRUE`:

```
ndvuong(nl, hl, vartest = TRUE)
```

```
variance test
```

```
data: nl-hl
w2 = 0.047327, p-value < 2.2e-16
alternative hypothesis: positive variance
```

The null hypothesis that  $\omega^2 = 0$  is rejected. We can then proceed to the second step, which is the test for non-nested models.

```
ndvuong(hl, nl)
```

```
Non-degenerate Vuong test for non-nested models
```

```
data: hl-nl
z = 1.7298, vuong_stat = 1.829208, constant = 0.000000, sum e.v. =
-1.832021, vuong_p.value = 0.033684, p-value = 0.0975
alternative hypothesis: different models
```

The classical and the non-degenerate Vuong tests both conclude that the two models are equivalent at the 5% level, but that the heteroscedastic model is better than the nested logit model at the 10% level.

## References

Bhat, Chandra R. 1995. “A Heteroscedastic Extreme Value Model of Intercity Travel Mode Choice.” *Transportation Research Part B: Methodological* 29 (6): 471–83. <https://www.sciencedirect.com/science/article/pii/0191261595000156>.

- Coate, Stephen, and Michael Conlin. 2004. "A Group Rule-Utilitarian Approach to Voter Turnout: Theory and Evidence." *American Economic Review* 94 (5): 1476–1504.
- Croissant, Yves. 2020. "Estimation of Random Utility Models in r: The Mlogit Package." *Journal of Statistical Software* 95 (11): 1–41. <https://doi.org/10.18637/jss.v095.i11>.
- Koppelman, Frank S., and Chieh-Hua Wen. 2000. "The Paired Combinatorial Logit Model: Properties, Estimation and Application." *Transportation Research Part B: Methodological* 34 (2): 75–89. <https://www.sciencedirect.com/science/article/pii/S0191261599000120>.
- León, Gianmarco, and Edward Miguel. 2017. "Risky Transportation Choices and the Value of a Statistical Life." *American Economic Journal: Applied Economics* 9 (1): 202–28. <https://doi.org/10.1257/app.20160140>.
- Shi, Xiaoxia. 2015. "A Nondegenerate Vuong Test." *Quantitative Economics*, 85–121.
- Vuong, Quang H. 1989. "Likelihood Ratio Tests for Selection and Non-Nested Hypotheses." *Econometrica* 57 (2): 397–33.
- Wen, Chieh-Hua, and Frank S Koppelman. 2001. "The Generalized Nested Logit Model." *Transportation Research Part B: Methodological* 35 (7): 627–41. <https://www.sciencedirect.com/science/article/pii/S019126150000045X>.