

Disposition of Participant
All Participants as Treated

	Placebo		Low Dose		High Dose		Total	
	n	(%)	n	(%)	n	(%)	n	(%)
Participants in population	86		84		84		254	
Trial Disposition								
Complete	47	(54.7)	34	(40.5)	38	(45.2)	119	(46.9)
Discontinued	26	(30.2)	37	(44.0)	38	(45.2)	101	(39.8)
Lack of Efficacy	2	(2.3)	3	(3.6)	9	(10.7)	14	(5.5)
Lost to Follow-Up	2	(2.3)	7	(8.3)	8	(9.5)	17	(6.7)
Other	15	(17.4)	18	(21.4)	18	(21.4)	51	(20.1)
Withdrawal By Subject	7	(8.1)	9	(10.7)	3	(3.6)	19	(7.5)
Participants Ongoing	13	(15.1)	13	(15.5)	8	(9.5)	34	(13.4)
Participant Study Medication Disposition								
Complete	47	(54.7)	34	(40.5)	38	(45.2)	119	(46.9)
Discontinued	26	(30.2)	37	(44.0)	38	(45.2)	101	(39.8)
Adverse Event	15	(17.4)	18	(21.4)	18	(21.4)	51	(20.1)
Lack of Efficacy	2	(2.3)	3	(3.6)	9	(10.7)	14	(5.5)
Lost to Follow-Up	2	(2.3)	7	(8.3)	8	(9.5)	17	(6.7)
Withdrawal By Subject	7	(8.1)	9	(10.7)	3	(3.6)	19	(7.5)
Participants Ongoing	13	(15.1)	13	(15.5)	8	(9.5)	34	(13.4)

Source: [CDISCpilot: adam-ads]